



A111.E1040  
JACC March 9, 2010  
Volume 55, issue 10A



## MYOCARDIAL ISCHEMIA AND INFARCTION

### REGIONAL BRAIN OXYGEN SATURATION AS A NOVEL INDEX OF NEUROLOGICAL OUTCOMES IN PATIENTS WITH OUT-OF-HOSPITAL CARDIOGENIC CARDIAC ARREST

ACC Poster Contributions

Georgia World Congress Center, Hall B5

Sunday, March 14, 2010, 3:30 p.m.-4:30 p.m.

Session Title: Pre-Hospital Emergency Therapies for Cardiac Arrest

Abstract Category: Cardiopulmonary Resuscitation/Emergency Cardiac Care/Shock

Presentation Number: 1106-314

Authors: *Noritoshi Ito, Shinsuke Nanto, Yasuji Doi, Hirotaka Sawano, Daisaku Masuda, Shizuya Yamashita, Mai Hatano, Daisuke Tonomura, Yuma Kurozumi, Tomoaki Natsukawa, Yusuke Ito, Kazuyuki Oka, Jiro Ooba, Taizo Hasegawa, Makoto Kobayashi, Hiroshi Ichiyanagi, Koji Akashi, Koichi Otsuya, Shoji Kaibe, Ken-ichiro Okada, Yasuyuki Hayashi, Tatsuro Kai, Toru Hayashi, Osaka Saiseikai Senri Hospital, Suita, Japan, Osaka University Graduate School of Medicine, Suita, Japan*

**Background:** Although ammonia (NH<sub>3</sub>) is a useful marker of brain resuscitation in patients with out-of-hospital cardiac arrest (OHCA), it takes a certain amount of time to measure the NH<sub>3</sub> level. If an objective index enabled the earlier prediction of a patient's outcome, it would help to determine whether more aggressive treatment should be performed on the patient. So we focused on noninvasive regional brain oxygen saturation (rSO<sub>2</sub>), because rSO<sub>2</sub> can be measured sooner than NH<sub>3</sub> by INVOS5100C (Somanetics, USA) and monitoring rSO<sub>2</sub> levels during cardiovascular surgery has been shown to relate to the neurological outcome. We investigated whether rSO<sub>2</sub> can be used as a useful index for estimating the neurological prognosis in patients with OHCA. To our knowledge, this is the first report on rSO<sub>2</sub> levels in patients with OHCA.

**Methods:** Of the 46 patients with OHCA during the period from March to August 2009, 25 of them were cardiogenic arrest. In these 25 patients, we statistically analyzed the correlation between the NH<sub>3</sub> levels and the rSO<sub>2</sub> levels measured immediately after arrival.

**Results:** The rSO<sub>2</sub> levels in 13 patients with the favorable neurological outcome were significantly higher than those in 12 patients with the unfavorable neurological outcome (a median, 60% vs 16%,  $p < 0.001$ ). There was a close negative correlation between the NH<sub>3</sub> levels and the rSO<sub>2</sub> levels.

**Conclusion:** rSO<sub>2</sub> is an easily obtained novel index for estimating the neurological prognosis in patients with out-of-hospital cardiogenic cardiac arrest.

**Figure:** Correlation between the NH<sub>3</sub> levels and the rSO<sub>2</sub> levels

